



Traumatic Injuries to Primary Teeth



م. هبة نافع ياسين
2022



Introduction

- ▶ Traumatic injuries to the primary dentition present special problems and the management is often different as compared with the permanent dentition.
- ▶ Trauma to the oral region occurs frequently and comprises 5% of all injuries for which people seek treatment .





Epidemiology of Dental Trauma



At 5 years, 31-40 % boys and 16-30 % girls will have suffered some dental trauma.

• At 12 years, 12-33 % boys and 4-19 girls will have suffered some dental trauma.

• Boys affected twice as often as girls.

• Maxillary anterior are the most affected teeth.


• Concussions, subluxation and luxation are the commonest injuries in primary dentition.





Etiology of Dental Trauma

- In primary teeth: Incompletely developed coordination leading to fall in and around home.
- In permanent teeth: Falls and collisions during outdoor activities.
- Examples: Sports injuries, road traffic accidents, child abuse
- Types: Direct trauma, indirect trauma



Predisposing Factors

- Protruded upper incisors
- Insufficient lip closure

1-Injuries to the Periodontal Tissues




- Concussion
- Subluxation



- Extrusive luxation
- Lateral luxation



- Intrusive luxation
- Avulsion



Clip slide


Injuries to The tooth

- Enamel infarction
- Enamel fracture
- Enamel – dentine fracture
- Complicated crown fracture
- Uncomplicated crown – root fracture
- Complicated crown root fracture
- Root fracture




Injuries to Gingiva and Oral Mucosa

- Laceration
- Contusion
- Abrasion



Injuries to the Supporting Bone




- **Comminution of socket wall**
- **Fracture of socket wall**
- **Fracture of alveolar process**
- **Fracture of jaw bones**



Treatment principle



The primary goal is to optimize periodontal and pulpal healing in the primary dentition provided that no further injury is transmitted to the developing permanent successors.





Dental And Medical History

- When, where, how?
- Lost teeth fragments?
- Headache, vomiting, amnesia?
- Bleeding disorders?
- Drug allergies?
- Tetanus immunization status?
- Rheumatic fever?



A photograph of a woman in a dental chair, looking upwards with her hands covering her mouth. A dentist in a white coat and blue gloves is examining her teeth. The background is a clinical setting with blue and white walls.

Extraoral Examination

- **General examination (look for signs of shock e.g. pallor, cold skin, irregular pulse, hypotension)**
- **Facial swelling, bruises, lacerations**
- **Limitation of mandibular movement or mandibular deviation**

Intraoral Examination



- Inspection (Laceration, abrasion, lip swelling, occlusal abnormality, fractured or cracked crowns, tooth color change)

- Palpation (mobility of tooth or alveolar process)

- Percussion (dull note?)

Treatment Of Traumatized Primary Teeth

Uncomplicated crown fracture	Enameloplasty, restoration
Complicated crown fracture	Pulpectomy, obturation with ZnO/ Eugenol; extraction
Crown – root fracture	Extraction
Root fracture	Remove coronal necrotic fragment
Concussion	Occlusal relief, analgesics
Subluxation	Soft diet, follow up; extraction
Extrusive luxation	Extraction
Intrusive luxation	Palatal displacement → extraction; buccal displacement → followup for spontaneous reeruption
Avulsion	No treatment (No replantation, no space maintainer)

خذ أول حرف من كل حيوان



إذا عرفت الجواب فأنت عبقرى

A young child with dark hair is crying in pain, holding their mouth with both hands. They are wearing a blue and white striped shirt. In front of them on a white table is a plate of colorful candies. The background is white with decorative geometric shapes: a pink semi-circle at the top, a yellow square on the right, a light blue plus sign in the lower right, and a dark blue diagonal bar at the bottom right. A vertical bar of colorful squares is on the far left edge.

Sequelae of Acute Dental Trauma in The Primary Dentition.

Pulpitis



Pulpitis is the initial response of the tooth to trauma and it accompanies almost every injury. Signs include sensitivity to percussion and capillary congestion, which may be clinically apparent from the lingual surface of the tooth using transillumination. Pulpitis may be reversible in minor injuries or may progress to irreversible pulpitis and pulp necrosis

Tooth Discoloration



- Injuries to the primary incisors frequently cause tooth discoloration
- **Blood vessels within the pulp chamber can rupture, depositing blood pigment in the dentinal tubules.**
- This blood may desorb completely or can persist to some degree throughout the life of the tooth

Crown Discoloration



- **Discoloration is a common complication after luxation injuries.**
- **Injured pulps may lose their vitality either because of damage to the vascular tissue at the apex and the resulting ischemia or because of necrosis of exposed coronal pulp tissue.**
- **If the necrotic pulp becomes infected with oral microorganisms either because of luxation of the root and ingress through the lacerated PDL or via an exposed pulp, pain and root resorption can occur**
- **Unless associated infection exists, root canal treatment is not indicated**



Teeth that discolor are not necessarily necrotic, particularly when the color change occurs within a few days of the injury.

A yellowish discoloration of both primary and permanent teeth may occur if they undergo pulp canal obliteration



Pulp Canal Obliteration

The entire pulp chamber and canal appear radiopaque in radiographs and the crown may have a yellowish color.



A young child with dark hair is shown in profile, crying with their mouth wide open and eyes closed. They are holding their right hand to their cheek. The child is wearing a blue and white striped t-shirt. The background is a plain, light-colored wall. On the left side of the image, there is a vertical bar with a series of colored squares in shades of red, blue, yellow, and orange. A small red horizontal bar is positioned above the child's head on the left side.

Sequelae in Permanent Dentition after Trauma in Primary Dentition.

👁 Enamel hypoplasia

- ✓ This includes discoloration of the enamel and or defects of the enamel surface.
- ✓ Discoloration usually ranges from white to yellowish-brown staining.
- ✓ The hypoplasia normally affects the labial crown surface and ranges from tiny spots to large areas.






Crown Dilaceration

- ✓ A traumatic displacement of already formed hard tooth substance in relation to the developing soft tissues leads to a deviation of the crown in relation to the long axis of the tooth.
- ✓ A minor dilaceration consists of a circular enamel defect.
- ✓ The severe type includes a complete palatal deviation of the crown with additional enamel hypoplasia

Odontome-like teeth



- ▶ Heavy trauma to the permanent tooth germ at an early stage of odontogenesis may lead to complete tooth deformation.
- ▶ Odontome-like disturbances of permanent teeth may develop especially after intrusive or luxation of primary teeth.
- ▶ On radiographs such malformed teeth present as a conglomeration of hard tissues resembling a complex odontome.
- ▶ As a rule such malformed teeth do not erupt and must be removed surgically.




Root Malformation

- ▶ Trauma to the epithelial root sheath of Hertwig during root development may lead to root dilaceration or to an arrest of root formation
- ▶ In the latter case a very short root may develop and tooth eruption will be delayed or completely disturbed.
- ▶ Other, but very rare, malformations include root duplication and lateral or vestibular root angulation



References

- ▶ Andersson L, Andreasen JO, Day P. Guidelines for the Management of Traumatic Dental Injuries: 2. Avulsion of Permanent Teeth. *Dent Traumatol* 2012;28:88-96.
 - ▶ Malmgren B, Andreasen JO, Flores MT, Robertson A, DiAngelis AJ, Andersson L, Cavalleri G, Cohenca N, Day P, Hicks ML, Malmgren O. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 3. Injuries in the primary dentition. *Dental Traumatology*. 2012 Jun 1;28(3):174-82
 - ▶ Holan G. Long-term effect of different treatment modalities for traumatized primary incisors presenting dark coronal discoloration with no other signs of injury. *Dental Traumatology*. 2006 Feb 1;22(1):14-7
- 



Thank you

م. هبة نافع ياسين

جامعة بغداد - كلية طب الاسنان

طب اسنان الاطفال والوقائي

2022

